

The Greenland Eskimos craniometric diversity

Elżbieta Jaskulska

Department of Historical Anthropology
Institute of Archaeology, Warsaw University

Keywords: Eskimo, Greenland, metric diversity

Introduction

This paper presents morphological analysis of Greenland Eskimos skulls. It is widely accepted that Greenland inhabitants, living in one of the most inhospitable areas people have ever populated, constitute an interesting group for research on ecologically based differentiation of *Homo sapiens* (e.g. Koertvelyessy 1972, Ruff 1994). Craniometric characteristics of human morphology have been for long time one of the main aims of arctic anthropology (Hrdlička 1910, Furst and Hansen 1915, Oschinsky 1962, Zegura 1978), and as Eskimo groups have a long history of being researched anthropologically, they consist one of the most commonly used reference group for analysing climatic differentiation of human species. Most researchers treat Eskimos as morphologically homogenous, distinguishing at best four main group: Asian, South Alaskan, Canadian and Greenland Eskimos. Although this distinction is well founded on the cultural and ethnological basis (Dumond 1987) some researchers give also indication for further division.

The main advantage of the population of Greenland Eskimos is a relatively simple model of settlement on the island, with habitable area occurring only on the shores and clear migration routes (Dumond 1987, see also Figure 1). This provokes some authors to state that the groups on the ends of different migration routes (i.e. NE i SE groups) can be treated as isolates (Laughlin 1966). If such opinion is well grounded by anthropological research, it can give some insight into process of adaptation to the climate of the group living in extremely harsh environment.

Table 1: List of samples with names of archaeological sites and their locations

| <i>Site code</i> | <i>Location</i> | <i>Site name</i> | <i>Number of individuals</i> |
|------------------|-----------------|-------------------------------|------------------------------|
| 111 | NW | York Peninsula | 6 |
| 112 | NW | Upernivik (Upernavik) | 34 |
| 113 | NW | Umanak (Umânaq) | 38 |
| 114 | NW | Ritenbaenk & Disko | 35 |
| 115 | NW | Jacobshavn (Ilulissat) | 15 |
| 116 | NW | Christianshåb (Qasigiannguit) | 15 |
| 117 | NW | Egedesminde (Ausiait) | 20 |
| 121 | SW | Sukkertoppen (Manitsôq) | 10 |
| 122 | SW | Godthåb (Nûk) | 16 |
| 123 | SW | Frederikshåb (Pâmiut) | 11 |
| 124 | SW | Julianehåb (Qaqortoq) | 11 |
| 211 | NE | Steward Peninsula | 15 |
| 212 | NE | Storm Bay | 4 |
| 221 | SE | Kutek | 11 |
| 222 | SE | Angmagssalik (Ammassadik) | 13 |
| 223 | SE | Nualik | 17 |

Table 2: F-matrix between researched Greenland sample and Howells' data (only samples with distance smaller than Norse sample are presented, mean calculated from all distances to Howells' samples)

| <i>Sample</i> | <i>Location</i> | <i>Between groups F-matrix distance to 'x'</i> |
|-------------------------|-----------------------------|--|
| Eskimo | Greenland; America | 16.4800 |
| Zalavar | Hungary; Central Europe | 71.8680 |
| Guam | Oceania | 80.1310 |
| Atayal | Taiwan; Asia | 83.4830 |
| Anyang (Shang dynasty) | China; Asia | 90.0160 |
| Hokkaido | North Japan; Asia | 93.8750 |
| Santa Cruz Island | California; North America | 94.2470 |
| North Kyushu | South Japan; Asia | 99.9900 |
| Ainu | North Japan; Asia | 106.9520 |
| Norse (medieval) | Norway; North Europe | 112.9650 |
| MEAN | | 136.1685 |




Figure 1: Location of sites and migration routes

Materials and methods

Data used for present analysis comes from two comprehensive anthropological studies: Aleš Hrdlička’s “Catalog of human crania in United States National Museum Collections: Eskimo in general” (1942) and Furst and Hansen’s “Crania Groenlandica” (1915). There were overall 416 skulls originating from Greenland and 271 (123 females and 148 males) presenting individuals derived from specified locations were chosen for present analysis. The list of the sites is presented in Table 1 and their location on the Figure 1.

The individuals from both sources can be described as representing indigenous inhabitants of Greenland, but actual chronology of the samples remains unknown, though it can be safely assumed that majority of them represent Thule culture period (1000–1600 AD), as the remains of earlier inhabitants are scarcer (Oschinsky 1962). The data was checked for its uniformity, as unknown chronology of the researched samples can indicate some Caucasian influence. As a reference the Howells’ data were used (Howells 1989). F-matrix between groups from all over the world and researched Greenland sample were measured ($df_1=9$ $df_2=2787$). Calculated values are proportional to distance measures and are computed from Mahalanobis D^2 statistics. The smallest difference (distance 16.48 with mean distance of 136.1685, see also Table 2) occurred between researched sample (‘x’) and reference Eskimo group which consists of Green-

Table 3: Jackknifed classification matrix for researched Greenland Eskimos ('x') and Howells' Eskimos samples

| | 'x' | Eskimo | 'x' + Eskimo |
|-------------------------|--------|--------|--------------|
| 'x' | 184 | 24 | 208 |
| | 67.90% | 22.22% | 54.88% |
| Eskimo | 59 | 58 | 117 |
| | 21.77% | 53.70% | 30.87% |
| Santa Cruz Island | 6 | 6 | 12 |
| | 2.21% | 5.56% | 3.17% |
| North Kyushu | 3 | 4 | 7 |
| | 1.11% | 3.70% | 1.85% |
| Tolai | 3 | 4 | 7 |
| | 1.11% | 3.70% | 1.85% |
| Lake Alexandrina Tribes | 2 | 2 | 4 |
| | 0.74% | 1.85% | 1.06% |
| Moriori | 1 | 3 | 4 |
| | 0.37% | 2.78% | 1.06% |
| Ainu | 1 | 2 | 3 |
| | 0.37% | 1.85% | 0.79% |
| Egypt | 3 | | 3 |
| | 1.11% | | 0.79% |
| Norse (medieval) | 2 | 1 | 3 |
| | 0.74% | 0.93% | 0.79% |
| Yauyos | 2 | 1 | 3 |
| | 0.74% | 0.93% | 0.79% |
| Zalavar | 2 | 1 | 3 |
| | 0.74% | 0.93% | 0.79% |
| Arikara (early) | 1 | | 1 |
| | 0.37% | | 0.26% |
| Berg | 1 | | 1 |
| | 0.37% | | 0.26% |
| Easter Island | | 1 | 1 |
| | | 0.93% | 0.26% |
| Guam | | 1 | 1 |
| | | 0.93% | 0.26% |
| Zulu | 1 | | 1 |
| | 0.37% | | 0.26% |
| Total number | 271 | 108 | 379 |

land Eskimo without Caucasian admixture. Also discriminant analysis shows that only 2 of total number of 271 researched Eskimo skulls group together with Norse sample (also one of the original Howells' Eskimo are grouped together with Norse sample) – see jackknifed classification matrix for 'x' and Howells' Eskimo samples in Table 3.

Only most common cranial measurements that simultaneously well describe skull morphology were chosen. Analysed features were maximal cranial length (*gonion-opisthocranion*; M1 acc. to R. Martin 1957-66), (*eurion-aurion*; M8), cranial height (*basion-bregma*; M17), upper facial height (*nasion-prosthion*; M48), bizygomatic facial breadth (*zygion-zygion*; M45), nasal height (*nasion-nasospinale*; M55), nasal breadth (*aperthion-aperthion*; M54), orbital height (*subconchion-supraconchion*; M52) and orbital breadth (*dacryon-ectoconchion*; M51a) (see also Table 4). Only skulls with full data, together with precise information on their origins were taken into analysis. Table 5 presents mean values of all variables for each sample.

Table 4: List of taken measurements

| <i>Measurement</i> | <i>Points</i> | <i>acc. to Martin</i> | <i>acc. to Howells</i> |
|----------------------------|---------------|-----------------------|------------------------|
| maximal cranial length | g-op | M1 | GOL |
| maximal cranial breadth | eu-eu | M8 | XCB |
| cranial height | ba-b | M17 | BBH |
| upper facial height | n-pr | M48 | NPH |
| bizygomatic facial breadth | zy-zy | M45 | ZYB |
| nasal height | n-ns | M55 | NLH |
| nasal breadth | apt-apt | M54 | NLB |
| orbital height | sbk-spa | M52 | OBH |
| orbital breadth | d-ek | M51a | OBH |

Results

The data, as expected, can describe sexual dimorphism in skull size for the Greenland Eskimo, as t-test shows significant differences in all measurements. To exclude its influence on the results of the present analysis all data was standardised for each sex separately.

Analysis of single measurements shows some significant differences in variance between samples: in maximal cranial length, maximal cranial breadth and orbital breadth (see Table 6).

The data shows that all the skulls belong to one population without much morphological differences. When analysing matrix of the Mahalanobis D^2 distances between the samples, it can be easily seen that major distances occur between sample 111 from York Peninsula (Thule settlement¹), with maximal distance of 4.185 (underlined in Table 7) and highest mean distance from other samples equal to 2.4. This can be explained by ethnic difference of the Eskimos from that area: ethnographers classify inhabitants of Thule settlement and their closest neighbours as belonging to Polar Eskimo group, consisting of Eskimos of High Arctic Canadian Islands. So even the morphological differences are not strong enough to show significant results in variance analysis some morphological difference between this sample and rest of the Greenland population can be found. Closer analysis show that the differences in cranial length and orbital breadth between this sample and other is responsible for results: exclusion of 111 sample from ANOVA analyses shows insignificant difference in variance in cranial length ($f=1.725$, $p>0.05$) and orbital breadth ($f=1.621$, $p>0.073$).

Analysing the Mahalanobis distances between samples from East Greenland (bolded in Table 8) it can be easily seen that none of the distances between them occurs bigger than average distance between sample 111 and the rest of the Greenland inhabitants (mean distance = 2.4003). In fact mean distance between East Greenland samples (mean = 1.0525) are smaller than mean distances between all samples (mean = 1.498).

Checking the distance between the samples shows that similar morphological characteristics appear on the both shores of Greenland and there is no latitudinal

¹Thule settlement was an eponymous site for Thule culture.

Table 5: Mean values and Standard Deviation of all variables for every sample researched

| | GOL | XCB | BBH | NPH | ZYB | NLH | NLB | OBH | ORB |
|-----|--------------|--------------|--------------|-------------|--------------|-------------|-------------|-------------|-------------|
| 111 | 183.4 ± 6.11 | 134.4 ± 4.61 | 136.3 ± 4.39 | 70.7 ± 4.89 | 138.4 ± 7.93 | 53.4 ± 3.10 | 21.9 ± 1.68 | 36.0 ± 0.58 | 44.1 ± 2.04 |
| 112 | 186.2 ± 6.39 | 131.9 ± 4.60 | 137.3 ± 4.75 | 72.8 ± 4.88 | 136.3 ± 7.13 | 52.0 ± 3.82 | 23.0 ± 1.75 | 36.1 ± 1.59 | 41.9 ± 2.43 |
| 113 | 186.8 ± 6.55 | 133.2 ± 3.99 | 136.2 ± 4.78 | 73.7 ± 4.20 | 136.7 ± 7.00 | 52.5 ± 2.93 | 22.7 ± 1.66 | 35.9 ± 2.26 | 42.1 ± 3.01 |
| 114 | 187.1 ± 6.85 | 131.9 ± 5.30 | 137.6 ± 4.47 | 72.6 ± 4.88 | 136.3 ± 8.06 | 52.1 ± 3.33 | 23.3 ± 1.90 | 35.7 ± 2.31 | 42.5 ± 2.85 |
| 115 | 183.6 ± 5.74 | 134.3 ± 5.13 | 135.5 ± 4.59 | 72.1 ± 5.47 | 135.6 ± 5.90 | 52.5 ± 3.28 | 22.6 ± 2.34 | 35.4 ± 1.91 | 41.8 ± 3.07 |
| 116 | 188.0 ± 5.25 | 135.4 ± 3.79 | 136.1 ± 4.43 | 72.6 ± 5.12 | 136.8 ± 6.41 | 52.9 ± 4.66 | 22.6 ± 1.45 | 36.3 ± 1.69 | 41.7 ± 2.33 |
| 117 | 184.4 ± 7.54 | 133.8 ± 5.05 | 133.5 ± 5.73 | 73.0 ± 4.25 | 133.5 ± 7.69 | 51.9 ± 2.71 | 22.4 ± 1.97 | 35.9 ± 1.93 | 42.4 ± 2.22 |
| 121 | 182.9 ± 4.91 | 131.6 ± 4.46 | 134.5 ± 5.96 | 72.1 ± 3.11 | 133.7 ± 9.73 | 52.3 ± 2.65 | 22.4 ± 1.96 | 35.2 ± 1.25 | 42.6 ± 1.75 |
| 122 | 187.2 ± 7.00 | 132.2 ± 4.39 | 137.1 ± 5.23 | 72.6 ± 6.77 | 137.8 ± 7.43 | 52.8 ± 2.91 | 23.1 ± 2.27 | 35.8 ± 1.65 | 42.4 ± 2.33 |
| 123 | 185.6 ± 5.70 | 135.6 ± 3.67 | 138.1 ± 5.30 | 75.5 ± 3.45 | 137.5 ± 6.28 | 53.9 ± 1.58 | 22.0 ± 1.48 | 36.5 ± 2.91 | 42.9 ± 2.51 |
| 124 | 183.0 ± 4.67 | 133.8 ± 3.43 | 137.1 ± 4.37 | 73.3 ± 5.45 | 135.5 ± 6.19 | 53.2 ± 4.73 | 22.5 ± 1.88 | 34.8 ± 2.08 | 42.2 ± 1.60 |
| 211 | 183.9 ± 5.58 | 134.5 ± 5.50 | 134.3 ± 5.68 | 71.4 ± 4.99 | 134.4 ± 6.10 | 52.4 ± 2.99 | 21.9 ± 1.19 | 35.9 ± 2.24 | 41.6 ± 1.93 |
| 212 | 182.8 ± 4.86 | 135.5 ± 4.65 | 133.3 ± 4.92 | 74.0 ± 3.16 | 130.8 ± 4.99 | 52.8 ± 2.99 | 21.5 ± 1.73 | 35.0 ± 0.82 | 40.3 ± 1.89 |
| 221 | 182.9 ± 6.06 | 133.1 ± 5.19 | 135.7 ± 8.24 | 70.3 ± 4.41 | 135.8 ± 7.87 | 49.4 ± 3.23 | 22.3 ± 1.35 | 35.6 ± 2.16 | 40.0 ± 1.41 |
| 222 | 186.7 ± 6.10 | 131.0 ± 6.58 | 137.0 ± 4.83 | 74.8 ± 3.60 | 135.1 ± 7.34 | 53.4 ± 2.60 | 22.6 ± 1.89 | 35.0 ± 1.96 | 41.5 ± 2.15 |
| 223 | 183.7 ± 7.32 | 131.7 ± 4.75 | 136.5 ± 6.24 | 72.3 ± 5.88 | 132.2 ± 9.11 | 52.0 ± 3.85 | 22.9 ± 1.49 | 35.4 ± 2.25 | 40.8 ± 1.82 |

Table 6: The ANOVA results for researched measurements

| <i>Variable</i> | | <i>f</i> | <i>p</i> |
|--------------------------------|------------------|--------------|--------------|
| maximal cranial length | g-op/GOL | 1.894 | 0.024 |
| maximal cranial breadth | eu-eu/XCB | 2.046 | 0.013 |
| cranial height | ba-b/BBH | 1.204 | 0.268 |
| upper facial height | n-pr/NPH | 0.835 | 0.638 |
| bizygomatic facial breadth | zy-zy/ZYB | 1.576 | 0.080 |
| nasal height | n-ns/NLH | 1.195 | 0.275 |
| nasal breadth | apt-apt/NLB | 1.142 | 0.319 |
| orbital height | sbk-spa/OBH | 0.787 | 0.692 |
| orbital breadth | d-ek/OBB | 1.766 | 0.039 |

differences, as it is presented in the cluster tree on the Figure 2. The clustering was based on the Mahalanobis distances, and was performed using average linkage method (farthest neighbour).

Discussion

The graph shows two main clusters and series from all part of Greenland appear in both clusters. This justifies the statement that Eskimo population in Greenland is rather homogenous apart of the maximal cranial breadth, which differs between series with no clear trend occurring (see Figure 3).

The only exclusion from this rule can be a York Peninsula settlement (sample 111), which differs significantly in maximal cranial length and orbital breadth.




Figure 2: Cluster tree based on Mahalanobis distances using average linkage method (farthest neighbour)

Table 7: The matrix of Mahalanobis distances between samples (distances between East Greenland samples are bolded)

| | 111 | 112 | 113 | 114 | 115 | 116 | 117 | 121 | 122 | 123 | 124 | 211 | 212 | 221 | 222 | 223 | mean |
|-----|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|-------|--------------|--------------|--------------|--------------|--------------|--------|
| 111 | 0.000 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 2.4003 |
| 112 | 3.568 | 0.000 | - | - | - | - | - | - | - | - | - | - | - | - | - | - | 1.5939 |
| 113 | 3.362 | 0.938 | 0.000 | - | - | - | - | - | - | - | - | - | - | - | - | - | 1.3112 |
| 114 | 3.384 | 0.691 | 1.660 | 0.000 | - | - | - | - | - | - | - | - | - | - | - | - | 1.9414 |
| 115 | 1.874 | 1.364 | 0.863 | 1.750 | 0.000 | - | - | - | - | - | - | - | - | - | - | - | 1.0039 |
| 116 | 2.697 | 1.461 | 1.004 | 1.890 | 0.873 | 0.000 | - | - | - | - | - | - | - | - | - | - | 1.4134 |
| 117 | 2.522 | 3.000 | 1.479 | 3.532 | 1.038 | 1.594 | 0.000 | - | - | - | - | - | - | - | - | - | 1.6680 |
| 121 | 1.032 | 1.330 | 0.773 | 1.336 | 0.496 | 1.100 | 0.597 | 0.000 | - | - | - | - | - | - | - | - | 0.9299 |
| 122 | 2.326 | 0.518 | 1.034 | 0.515 | 0.882 | 0.801 | 2.262 | 0.831 | 0.000 | - | - | - | - | - | - | - | 1.2821 |
| 123 | 1.698 | 2.205 | 1.393 | 2.752 | 0.832 | 1.713 | 0.801 | 0.575 | 2.115 | 0.000 | - | - | - | - | - | - | 1.3798 |
| 124 | 1.286 | 2.315 | 1.715 | 2.423 | 0.466 | 2.001 | 1.593 | 0.547 | 1.660 | 0.653 | 0.000 | - | - | - | - | - | 1.3931 |
| 211 | 1.777 | 2.154 | 1.183 | 2.724 | 0.604 | 0.382 | 1.259 | 0.632 | 1.251 | 0.955 | 1.143 | 0.000 | - | - | - | - | 1.3421 |
| 212 | 2.279 | 1.808 | 1.060 | 2.157 | 0.869 | 1.043 | 0.661 | 1.027 | 1.759 | 0.617 | 0.959 | 0.780 | 0.000 | - | - | - | 1.1821 |
| 221 | 4.185 | 1.679 | 1.861 | 3.092 | 1.737 | 2.187 | 2.701 | 2.278 | 2.165 | 2.492 | 2.639 | 2.473 | 1.690 | 0.000 | - | - | 2.1895 |
| 222 | 3.110 | 1.128 | 0.908 | 1.274 | 1.289 | 1.829 | 1.729 | 0.947 | 1.099 | 1.521 | 1.336 | 1.862 | 1.036 | 2.132 | 0.000 | - | 1.3645 |
| 223 | 3.304 | 1.344 | 1.746 | 1.882 | 1.126 | 2.039 | 1.920 | 1.378 | 1.296 | 1.755 | 1.553 | 2.294 | 1.168 | 1.721 | 0.632 | 0.000 | 1.5724 |




Figure 3: Results of ANOVA analysis of maximal cranial breadth in Greenland samples




Figure 4: Graphical presentations of Mahalanobis distances between samples from main parts of Greenland

Such statement stands in discrepancy with assumption that populations of different parts of Greenland, especially from NE and SE of Greenland are thought of as isolates (e.g. Laughlin 1966: 478). The present data shows that the most distant group of Greenland inhabitants is SE part (by Mahalanobis distances), whereas distance between NE and SE groups is only second biggest (see Table 8 and Figure 4).

Table 8: The Mahalanobis distances between samples from main parts of Greenland

| | NE | NW | SE | SW |
|----|-------|-------|-------|-------|
| NE | 0.000 | | | |
| NW | 1.906 | 0.000 | | |
| SE | 2.253 | 1.860 | 0.000 | |
| SW | 1.159 | 1.179 | 2.305 | 0.000 |

Table 9: T-test results for differences between groups. P<0.005 are bolded.

| variable | | t-test results | | | |
|----------------------------|-------------|----------------|--------|---------------|---------------|
| | | NE-SE | NW-SW | NW-NE | SW-SE |
| maximal cranial length | g-op/GOL | -0.413 | 1.062 | -1.570 | -0.471 |
| maximal cranial breadth | eu-eu/XCB | 1.970 | -0.100 | 1.415 | -1.353 |
| cranial height | ba-b/BBH | -1.435 | -0.533 | -1.920 | -0.234 |
| upper facial height | n-pr/NPH | -0.496 | -0.540 | -0.824 | -0.611 |
| bizygomatic facial breadth | zy-zy/ZYB | -0.157 | -0.255 | -1.417 | -1.447 |
| nasal height | n-ns/NLH | 0.823 | -1.367 | 0.245 | -1.849 |
| nasal breadth | apt-apt/NLB | -2.073 | 0.782 | -2.331 | 0.206 |
| orbital height | sbk-spa/OBH | 0.735 | 0.917 | -0.286 | -0.612 |
| orbital breadth | d-ek/OBB | 1.046 | -0.844 | -1.366 | -4.205 |

Closer analysis of differences (t-test) between NE and SE groups shows significant results for nasal breadth ($t=-2.073$; $df=59$; $p<0.05$), which suggests some isolation of both groups, which does not occur on the West Coast (see Table 9). It seems nonetheless that it is not enough to treat north-east and south-east groups as isolates. Analysis of differences between groups from east coast and adequate north-west and south-west groups shows that there are also significant differences between them (NW-NE also in nasal breadth: $t=-2.331$, SE-SW in orbital breadth: $t=-4.205$; for both $p<0.05$). It can be also noticed that those significant differences in nasal breadth are in both cases caused by wider than in other Greenland groups nasal apertures in the NE sample. None-the-less it has to be stated that this difference is not significant, as analysis of variance of nasal breadth between the samples shows insignificant results ($f=1.872$; $p>0.1$).

As those differences are bigger (SW-SE), or similar to NE-SE difference, assumption that only populations from east Greenland can be treated as isolates seem improbable on the ground of present research. Moreover the Greenland inhabitants form quite homogenous and morphologically similar population, which can be useful as a study group for climatic influence research.

Bibliography

Dumond D.E.

1987: *Eskimos and Aleuts*, London.

Furst C.M., Hansen F.C.C.

1915: *Crania Groenlandica. A Description of Greenland Eskimo Crania with an Introduction on the Geography and History of Greenland*, Copenhagen.

Howells W.W.

1989: *Skull Shapes and the Map. Craniometric Analyses in the Dispersion of Modern Homo*, Cambridge, Mass.

Hrdlička A.

1942: *Catalog of Human Crania in the United States National Museum Collections: Eskimo in General*, "Proceedings of the United States Museum", No. 3131, vol. 91, Washington, pp. 169-429.

Laughlin W.S.

1966: *Genetical and Anthropological Characteristics of Arctic Populations*, in: "The Biology of Human Adaptability", ed. Baker P.T., Weiner J.S., Oxford, pp. 469-508.

Martin R., Saller K.

1957: *Lehrbuch der Anthropologie in systematischer Darstellung mit besonderer Berücksichtigung der anthropologischen Methoden für Studierende, Ärzte und Forschungsreisende*, Stuttgart 1957-1966.

Oschinsky L.

1964: *The Most Ancient Eskimo. The Eskimo Affinities of Dorset Culture Skeletal Remains*, Ottawa.

Zegura S.L.

1978: *The Eskimo Population System: Linguistic Framework and Skeletal Remains*, in: "Eskimos of Northwestern Alaska. A biological Perspective", ed. Jamison P.L., Zegura S.L., Milan F.A., Stroudsburg, Pennsylvania, pp. 8-30.

